

Managing Patients with Feline Lower Urinary Tract Disease (FLUTD)/Urolithiasis

FELINE Lower Urinary Tract Disease (FLUTD)

Initially feed **Prescription Diet® c/d® Multicare Feline pet food** (e.g., during investigation)

Struvite

Urolith(s), urethral plug, or clinically significant crystalluria

Calcium oxalate, mixed struvite/calcium oxalate

Urolith(s) or clinically significant crystalluria

Struvite Dissolution

Feed **Prescription Diet® s/d® Feline pet food**

Dissolves struvite in as little as 8 days²

Not recommended for long-term feeding (no longer than 6 months)



Feed **Prescription Diet® c/d® Multicare Feline pet food**

Dissolves struvite uroliths in as little as 14 days¹

Ideal for long-term feeding



Remove uroliths
(e.g., surgery)

Aid in the management of occurrence or recurrence

Cats of ideal body weight

Feed **Prescription Diet® c/d® Multicare Feline pet food**



Obese prone cats

Feed **Prescription Diet® w/d® Feline pet food**



¹ Average 28 days. Lulich JP, et al. Struvite urolith dissolution in cats: a double blind randomized clinical trial of two foods *J Vet Intern Med* 2011;25:747.

² Average dissolution time 12 days. Data on file. Hill's Pet Nutrition, Inc.

Managing Patients with Feline Lower Urinary Tract Disease (FLUTD)/Urolithiasis

